

PROPAGATION OF *HEPTACODIUM MICONIOIDES*, SEVEN SON FLOWER, BY SOFTWOOD AND SEMI-HARDWOOD CUTTINGS

CHIN CHIN LEE AND T.E. BILDERBACK¹

*Department of Horticultural Science
North Carolina State University
Raleigh, North Carolina 27695-7609*

Abstract. Softwood and semi-hardwood two node cuttings of *Heptacodium miconioides* from basal and middlestem sections rooted better than terminal sections. Basal and middle softwood cuttings exhibited greater rooting (65 and 55%, respectively), than terminal cuttings (42%). Basal softwood cuttings produced more roots (7.0) than terminal cuttings (3.8). Root length and rootball diameter were not different among the three cutting positions. Semi-hardwood basal cuttings produced an average of 53.5 roots while middle and terminal cuttings produced 25.8 and 19.7, respectively. Cuttings treated with the potassium salt formulation of indolebutyric acid (K-IBA) exhibited increased rooting, greater root number and length, and greater rootball diameter in softwood cuttings. Semi-hardwood cuttings treated with K-IBA rooted in higher percentages and produced more roots than untreated cuttings.

INTRODUCTION

Heptacodium miconioides (Syn. *H. jasminoides* H.K.A. Shaur; cf Hu, 1988) is a small deciduous flowering tree in the Caprifoliaceae family, native to Western Hupeh, China (1). It is considered a rare plant even in its native habitat and was only available to the western world following the 1980 Sino-American Botanical Expedition to China (2,3). *Heptacodium miconioides* has potential as a new nursery crop because of its exfoliating bark, vigorous growth, and fragrant white late-summer flowers. After flowering, the persistent calyces develop an attractive reddish color in the fall.

Preliminary studies in 1987 by the authors compared single-node softwood cuttings taken from the terminal, middle and basal sections of stems. The basal portion of each cutting was immersed in a 50% ethanol solution containing one of three IBA treatments: 0, 2500, or 10,000 ppm indolebutyric acid (IBA). Results indicated basal sections rooted better than middle or terminal sections. Rooting for terminal, middle and basal sections was 33, 48, and 76%, respectively. However, many of the severed portions of the cuttings were necrotic. This may have been caused by the alcohol solvent or use of softwood cuttings.

Therefore, two studies were initiated with the objective of evaluating the rooting response of softwood and semi-hardwood

¹ Former graduate student, presently Lecturer, Department of Agronomy and Horticulture, University of Agriculture, Malaysia; and Associate Professor, respectively

cuttings from terminal, middle, and basal sections treated with selected concentrations of the potassium salt (K-salt) formulation of IBA (K-IBA).

MATERIALS AND METHODS

Experiment 1. Softwood cuttings were prepared on May 20, 1988 from 2-year-old container-grown *Heptacodium* plants maintained under natural daylength greenhouse conditions. The excised shoots had leaves that were fully expanded and light green in color. The stems were light green, and firm, and the immature distal stems were quadrangular in shape. Each cutting consisted of 2 nodes and was 8 to 10 cm (3.1 to 4.0 in) in length. A light wound was administered to each cutting consisting of 2 equidistant vertical incisions on the basal stem to a depth reaching secondary xylem, each about 3.0 cm (1.2 in) long. The basal 3.0 cm were then dipped for 10 sec. into the following concentrations of K-IBA: 0, 2,500, 5,000, and 10,000 ppm. After preparation, the basal end of 3.5 cm was inserted into a medium of perlite: peat (3:1 v/v). Each cutting was inserted into one of 24 individual cells in a plastic tray measuring 30 x 50 x 10 cm (11.8 x 19.7 x 3.9 in).

The experiment was conducted in a glass-covered greenhouse maintained at day/night temperatures of 30 and 20° C (86 and 68° F). Intermittent mist operated 3 sec every 5 min from 8:00 am. to 6:00 pm. daily. The experimental design was a randomized complete block design consisting of 12 trays and 24 replications. Each tray contained two completely randomized replications.

Twelve weeks after the experiment was initiated, data were recorded. Data included percent rooting, number and length of roots, and root-ball diameter. Any cutting with one or more roots 1 mm (0.04 in.) long was classified as rooted. Standard analysis of variance procedures were utilized for node position data analysis and regression for quantitative hormone concentration treatment analysis.

Experiment 2. On August 12, 1988, semi-hardwood cuttings were taken from the same stock plants utilized in Experiment 1, and divided into terminal, middle, and basal sections. The leaves were larger and darker green than the softwood cuttings and the stems were rounded with a reddish-green color. The semi-hardwood cuttings snapped with a distinct sound when broken. Two-nodal cuttings from each of the 3 sections were taken. Each cutting was 8 to 10 cm (3.1 to 4.0 in.) long. All cuttings were wounded as described in Experiment 1 and treated by dipping the basal 3.5 cm (1.4 in.) for 10 sec in one of the following concentrations of K-IBA: 0, 2,500, 5,000, 7,500, and 10,000 ppm. After air drying for approximately 8 min., cuttings were inserted

to a depth of 3.5 cm (1.4 in.) into individual cells in a rooting tray measuring 30 x 50 x 10 cm (11.8 x 19.7 x 3.9 in.).

Intermittent mist operated 3 sec every 5 min from 8:00 am. to 6:00 pm. daily. The experimental design was a randomized complete block using 4 cuttings per treatment with 6 replications. Each replication consisted of 2.5 trays for a total of 360 cuttings. The cuttings were evaluated after 12 weeks. Data included percent rooted, number, length of roots, and rootball diameter.

RESULTS AND DISCUSSION

Node position influenced percent rooting and root number in both experiments. However, there was no interaction between node position and rooting hormone concentrations, therefore data presented are averaged across nodal positions.

Experiment 1. Softwood cuttings from the basal and mid-sections rooted better than terminal cuttings (Table 1). The rooting percentage obtained was 64.6, 55.2, and 41.7%, respectively. Basal softwood cuttings had more primary roots than terminal cuttings (7.0 to 3.8). Primary root length and root-ball diameter were not different among the three nodal positions.

Table 1. Node position effect on rooting of softwood and semi-hardwood cuttings of *Heptacodium maconoides*²

Type of cutting	Rooting (%)	Root no.	Root length (mm)	Root-ball diam (mm)
<i>Experiment 1</i>		<u>Softwood</u>		
Basal	64.6 a	7.0 a	65.3 a	47.2 a
Mid-section	55.2 a	5.6 ab	60.7 a	41.1 a
Terminal	41.7 b	3.8 b	61.0 a	45.9 a
<i>Experiment 2</i>		<u>Semi-hardwood</u>		
Basal	85.0 a	53.5 a	87.6 a	61.8 a
Mid-section	60.8 b	25.8 b	78.2 a	41.9 b
Terminal	37.5 c	19.7 b	84.2 a	51.6 ab

² Means of 24 cuttings. Mean separation within column by LSD, 5% level

K-IBA application significantly increased rooting percentage, primary root number, and root-ball diameter in softwood cuttings (Table 2). The highest primary root number was obtained by treating cuttings with 5,000 ppm K-IBA. Root length was reduced when treated with 10,000 ppm K-IBA.

Table 2. Effect of K-IBA concentrations on rooting of softwood and semi-hardwood cuttings of *Heptacodium miconioides*.

K-IBA concentration	Rooting (%)	Root no	Root length (mm)	Root-ball diam. (mm)
<i>Experiment 1</i>		<u>Softwood</u>		
Treatment				
Nontreated	30.6	0.6	61.7	21.1
2,500 ppm	58.3	5.2	69.8	50.2
5,000 ppm	68.1	8.7	67.1	49.1
10,000 ppm	58.3	7.3	50.6	46.7
Significance				
R ²	0.11	0.14	0.05	0.12
Linear	***	***	NS	***
Quadratic	***	**	*	***
<i>Experiment 2</i>		<u>Semi-hardwood</u>		
Treatment				
Nontreated	30.6	14.7	95.4	45.3
2,500 ppm	65.3	34.6	86.4	50.9
5,000 ppm	76.4	42.1	84.3	55.5
7,500 ppm	70.8	40.8	76.7	53.3
10,000 ppm	62.5	32.8	77.6	52.8
Significance				
R ²	0.26	0.27	0.03	0.01
Linear	***	*	NS	NS
Quadratic	***	NS	NS	NS

NS, *, **, ***. Nonsignificant or significant at the 5%, 1%, or 0.1% levels, respectively

Experiment 2. Basal cuttings from semi-hardwood cuttings yielded a higher percent rooting, root number, and root-ball diameter than mid-section and terminal cuttings (Table 1). Basal cuttings produced more than twice as many primary roots as terminal cuttings (53.5 and 19.7 resp.). Node position did not influence root length.

Rooting percentage and primary root number were significantly increased with application of K-IBA in semi-hardwood cuttings ($P < 0.1$ and 0.05). A higher rooting percentage and a greater number of roots were produced with cuttings treated with K-IBA compared to nontreated cuttings of semi-hardwood materials. There were no K-IBA treatment effects on root length or root-ball diameter.

Data on growth after propagation is unavailable. *Heptacodium* in the North Carolina State Arboretum has reached a height of 3.6 m (12 ft.) and approximately half this width, but has been pruned extensively for propagation. Koller (2) reported that after five growing seasons seedling trees ranged from 1.8 to 3.0 m (6 to 10 ft.). Shoots from lateral cuttings appear to exhibit

plagiotrophic growth. However, they grow vigorously, and vertical shoots usually are initiated from the base of the plant.

LITERATURE CITED

1. Hu, J.Q. 1988 *Heptacodium*, in *Flora Republicae Popularis Sinicae*, Beijing Science Press. 72: 108-110
2. Koller, G L. 1986. Seven-son flower from Zhejiang. Introducing the versatile ornamental shrub, *Heptacodium miconioides* Airy Shaw. *Arnoldia* 46 (4):2-14
3. Koller, G.L. 1987 *Heptacodium miconioides*: A new small tree from China. *Amer. Nurs.* 165 (9).58-64